

FOR EXTERNAL ANIMAL USE ONLY

**CLOUT®**  
Reg. No. G1447 Act 36/1947  
N-SR0885

### POUR-ON FOR CATTLE, SHEEP AND GOATS

#### INDICATIONS:

##### CATTLE

- Controls ticks
- Controls stable flies, horn flies, cattle louse flies and nuisance flies e.g. house flies
- Kills lice (biting and sucking)
- Protects against blackflies

##### SHEEP AND GOATS

- Controls Karoo paralysis, bont-legged and red-legged ticks

**Contains:** Deltamethrin 1 % m/v

**DOSE:** Cattle - 1 ml/10 kg      Sheep and goats - 1 ml/5 kg

#### STORAGE INSTRUCTIONS

Store away from food and feed.

Tightly reseal the container if all the contents have not been used.

#### WARNINGS

- DO NOT USE ON CALVES/LAMBS UNDER 1 MONTH.
- NO WITHDRAWAL PERIOD FOR MEAT AND MILK.
- Avoid contact with product.
- Keep out of reach of uninformed persons, children and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.
- N.B. Pyrethroid pour-ons may cause irritation. Treat a small group of animals first and observe for an hour before treating the rest.

#### DIRECTIONS FOR USE: USE ONLY AS DIRECTED

##### Cattle

- **Ticks, flies and lice:** 1 ml/10 kg.

Apply along the backline, from the shoulder to the root of the tail using measuring dispenser or any other suitable applicator.

**In case of heavy brown ear or blue tick challenge**, apply Clout from the base of the skull to the root of the tail.

**Treat at regular intervals according to tick challenge.** The longer the product is regularly applied, the more the tick population is reduced. It will then become possible to apply at intervals greater than 7 days.

- **For blackfly (*Simulium*)** apply also to ears and underline. Repeat as required according to challenge.
- **For population control of nuisance flies and stable flies**, use 2 - 3 treatments at weekly intervals. Thereafter use according to challenge.

##### Sheep and goats

Dosage for optimal residual action – 1 ml/5 kg.

Environmental conditions and parasite challenge can influence treatment intervals.

- **Karoo paralysis ticks:** Divide the full dose per animal into 4 equal parts. Apply one part onto each axillae and groin region, whilst the animal is turned over in a sitting position.
  - Angora goats - repeat after approximately 2 weeks.
  - Dorper sheep - repeat after approximately 4 weeks.
  - Merino sheep - repeat after approximately 6 weeks.
- **Bont-legged and red-legged ticks:**

The dosage should be divided into 3 equal parts and applied to the axillae, groin and anogenital regions.

Treat sheep at regular intervals of approximately 3 - 4 weeks or as necessary, according to tick challenge.

For crippled sheep apply 1 ml to the affected hoof/fetlock.

**PRESENTATION: 200 ml, (500 ml, 1 l, 5l, 20l)**

#### Registration holder:

Cooper Veterinary Products (Pty) Ltd  
Co. Reg. No. 2002/021376/07  
P O Box 46, Isando 1600, RSA

#### Marketed by:

Afrivet Business Management (Pty) Ltd  
Co. Reg. No. 2000/011263/07  
P O Box 12406, Onderstepoort 0110, R.S.A.  
Tel: + 27 (0)12 545-0004