

FOR EXTERNAL ANIMAL USE ONLY

COOPERS REDLINE POUR-ON

Reg. No. G3445 Act 36/1947

READY-TO- USE POUR-ON FOR TICKS ON CATTLE, SHEEP, OSTRICHES AND GAME

- Controls ticks on cattle, sheep, ostriches and game.
- Controls Tsetse fly worry on cattle.
- Controls feather mite on ostriches.
- Sterilises female ticks.
- Easy application.
- Residual action.
- Non-systemic action.
- The product is rainfast.

Contains Flumethrin 1 % m/v

STORAGE INSTRUCTIONS

Protect from extremes of heat, cold and from dampness.

Store away from food and feed.

Tightly reseal the container if all the contents have not been used.

WARNINGS

- WITHDRAWAL PERIOD FOR MILK: NO WITHDRAWAL.
- WITHDRAWAL PERIOD FOR MEAT: 7 DAYS.
- Toxic to bees and fish.
- Although the colour of the product may vary, it will not influence the efficacy or residual period thereof.
- Skin reactions may occur in a small percentage of cases. If this happens, wash the affected animals with soap and water.
- Keep out of reach of uninformed persons, children and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Do not eat, drink or smoke during or after handling this product, before washing hands and face thoroughly.
- Avoid contact with pour-on and wash immediately and thoroughly with soap and hot water after accidental contamination.
- Remove and wash all contaminated clothing.
- Destroy empty containers and do not re-use for any other purpose.
- Do not contaminate dams, rivers and other water sources.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

- SHAKE WELL BEFORE USE.
- Treat at regular interval according to tick challenge.
- The longer the product is regularly applied, the more the tick population is reduced. It may then become possible to treat at extended intervals.

Dosage

The basic dosage is 1 ml per 10 kg body mass (refer to dosage table below).

- **Cattle:** Apply the required dose in two lines, from the base of the skull along both sides of the spine up to the tail root. For Tsetse fly treat every 14 days or according to challenge.
- **Sheep:** Apply the required dose volume in 4 parts to the bare skin areas of the groin and axillae. In the case of sheep infested with Karoo-paralysis tick (*Ixodes rubicundus*), Redline has to be applied early in the season when these ticks are first observed (either on the grazing or on the animals). Two to three treatments 7 to 10 days apart at this time should protect the sheep against Karoo-paralysis, depending on the severity of the tick challenge.
- **Game:** Apply the required dose volume with a suitable applicator. If self-medicating devices are used, it is important that the device only remains activated for as long as it takes to deliver one dosage to the animals in a particular paddock / camp. The device should not be used for continuous, low-level treatment.
- **Ostriches:** Apply the required dose volume along the backline.

Dosage table: Cattle and Game.

Body mass	Dosage	No. of doses per ℓ
Up to 100 kg	1 ml per 10 kg	
100 kg	10 ml	100
200 kg	20 ml	50
300 kg	30 ml	33
400 kg	40 ml	25
500 kg and over	50 ml	20

Dosage table: Sheep and Ostriches.

Body mass (kg)	Dose	No. of doses per ℓ
20	2 ml	500
30	3ml	333
40	4ml	250
50	5ml	200
60	6ml	166

Applicator

Squeeze bottle with dose meter. Squeeze required volume into dose meter, or use your drench gun.

200 ml (500 ml, 1ℓ, 5 ℓ, 6 ℓ, 20 ℓ)

Registration holder:

Cooper Veterinary Products (Pty) Ltd
Co. Reg. No. 2002/021376/07
P O Box 46, Isando 1600, RSA

Marketed by:

Afrivet Business Management (Pty) Ltd
Co. Reg. No. 2000/011263/07
P O Box 12406, Onderstepoort 0110, R.S.A.
Tel: + 27 (0)12 545-0004

