

FOR ANIMAL USE ONLY

OstriDose
Reg. No. G2234 Act 36/1947

A remedy against roundworms in ostriches and horses.

STORE IN A COOL, DRY PLACE

Composition: Fenbendazole 5 % m/v

Presentation: 1 l, 5 l, 10 l

WARNINGS

DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 7 DAYS OF LAST TREATMENT.

Do not mix or dilute with any other substance.

Keep out of reach of children, uninformed persons and animals.

Although this remedy has been extensively tested under a large range of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DIRECTIONS FOR USE – USE ONLY AS DIRECTED

Use as an oral drench.

Shake well before use.

Dosage:													
Ostriches:	3,0 ml /10 kg body mass (15,0 mg/kg)												
Horses:	2 ml /10 kg body mass (10,0 mg/kg)												
Example:													
Ostriches							Horses						
Body mass (kg)	Dose (3,0 ml /10 kg)	No. of doses in pack					Body mass (kg)	Dose (2,0 ml/10 kg)	No. of doses in pack				
		1 l	2 l	5 l	10 l	20 l			1 l	2 l	5 l	10 l	20 l
10	3	333	666	1666	3333	6666	100	20	50	100	250	500	1000
20	6	166	333	833	1666	3333	150	30	33	66	166	333	666
30	9	111	222	555	1111	2222	200	40	25	50	125	250	500
40	12	83	166	416	833	1666	250	50	20	40	100	200	400
50	15	66	133	333	666	1333	300	60	16	32	83	166	333
60	18	55	111	277	555	1111	350	70	14	28	71	142	285
70	21	47	95	238	476	952	400	80	12	24	62	125	250

EFFICACY

Ostriches:

* *Libyostrongylus douglassii*.

Horses:

* Bloodworms (large strongyles), redworms (small strongyles), pinworm (*Oxyuris*), ascarids and stomach worm (*Habronema majus*).

* Control = > 90 % effective

The list contains the more important endoparasite species. This anthelmintic may also be effective against other species. For more information consult your veterinarian.

Registration holder: ECO Animal Health Southern Africa (Pty) Ltd

Co. Reg. No. 92/00835/07

P O Box 12406, Onderstepoort, 0110

Tel: (012) 545-0004