

FOR EXTERNAL ANIMAL USE ONLY

**COOPERS SUPADIP**  
Reg. No. G3349 Act 36/1947

For the control of ectoparasites on cattle, horses, sheep, goats and dogs.

- Controls ticks, mange and lice on cattle.
- Eradicates lice and keds on sheep and goats.
- Controls ticks on sheep and goats.
- Protects sheep against maggot fly infestation.
- Controls ticks and fleas on dogs.
- Controls ticks on horses.

**Contains/Bevat:** Chlorfenvinphos/Chlorfenvinfos 30 % m/v

**STORAGE INSTRUCTIONS**

- Store in a cool, dry place.
- Store in original container with cap tightly closed.
- Store under lock and key.

**WARNINGS**

- DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION FOR 14 DAYS AFTER LAST TREATMENT.
- DO NOT USE ON LACTATING DAIRY COWS OR GOATS PRODUCING MILK FOR HUMAN CONSUMPTION.
- DO NOT DIP CALVES, LAMBS, KIDS OR FOALS UNDER 1 MONTH OF AGE.
- DO NOT DIP PUPPIES UNDER 3 MONTHS OF AGE, OR NURSING BITCHES.
- DO NOT DIP HOT, TIRED OR THIRSTY ANIMALS.
- DO NOT USE ON CATS OR PIGS.
- Toxic to fish – do not contaminate waterways.
- Toxic to insects, birds and other animals.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

**PRECAUTIONS**

- Do not smoke, eat or drink while handling this product.
- Wear protective clothing and rubber gloves.
- Avoid inhalation of spray wash.
- Wash immediately in the event of skin contact.
- Remove contaminated clothing and wash before re-use.
- Empty containers should be rinsed, punctured and disposed of in a safe manner.

**FIRST AID**

**Supadip is an organophosphate compound and toxic.**

**Symptoms of poisoning:**

Salivation, abdominal pain, diarrhea, muscle tremors and constriction of pupils.

Should the above symptoms appear seek medical advice and make this label available.

**Antidote:**

Atropine: 2 PAM

**DIRECTIONS FOR USE: USE ONLY AS DIRECTED**

Species	Method	Filling rate	Replenishment rate	Comments
Cattle	Dipping	1 : 600	1 : 320	Precream with an equal volume of water prior to addition to dip.
	Spraying	1 : 600	Boost rate: 185 ml Supadip/100 head sprayed	Make up to 400 l plus 2,5 l for every head to be sprayed.
	<b>Treatment intervals:</b> For routine tick control dip or spray weekly.			
Sheep	Dipping	1 : 600	1 : 300 when dip volume has fallen by 15 % to maintain strength.	OR use a constant replenishment system using 1 : 300 solution.
	Belly bathing	1 : 600	1 : 300 – Constantly replenish to maintain a depth of 60 cm.	Wet tails of sheep by submerging hindquarters as they walk through.
	Hand dressing	1 : 200 (50 ml Supadip in 10 l water)		For blowfly control: Apply to crutch and other common blowfly strike areas before the first rains and when necessary thereafter.
Goats	Dipping	1 : 600	1 : 300 when dip volume has fallen by 15 %.	For control of ticks.
		1 : 1200	1 : 600 when dip volume has fallen by 15 %.	For control of biting and sucking lice.
Sheep/Goats	<b>Treatment intervals:</b> <ul style="list-style-type: none"> <li>• Routine tick control – dip weekly.</li> <li>• Control of lice and keds – dip once only.</li> <li>• Control of blowfly strike – dip 4 – 6 weeks after shearing and repeat every 2 months.</li> </ul> <b>Note:</b> Where fouled dipwash is kept overnight for use the following day, add Copper Sulphate to prevent post dipping lameness (mix up to 2 kg Copper Sulphate in 5 l hot water and add at a rate of 1 : 800).			
Horses	Handspray /Hand dress	1 : 600		Thoroughly saturate coat. Pay particular attention to legs, belly and head.

	<b>Treatment intervals:</b> For tick control treat weekly.			
Dogs	Dog dip tanks	1 : 600	1 : 320 when dip volume has fallen by 15 %	Drain dip when wash becomes dirty to prevent unpleasant odour.
	Hand dressing	1 : 320 (10 ml/3,2 l water)		Mix required amount of solution. Thoroughly soak the coat and air or towel dry. DO NOT RINSE.
	<b>Treatment intervals:</b> <ul style="list-style-type: none"> <li>• For tick and flea control dip weekly in summer and every 2 weeks in winter. It is advisable to treat the dog's environment to control immature stages of fleas and ticks. Use remaining wash or a flea powder.</li> </ul> <b>NOTE:</b> Do not apply to puppies under 3 months of age or nursing bitches.			

- NOTE: DO NOT APPLY TO LACTATING DIARY COWS OR GOATS PRODUCING MILK FOR HUMAN CONSUMPTION.

Presentation: 100 ml, 250 ml, 500 ml, 1 l, 5 l, 10 l

**Registration holder:**

Cooper Veterinary Products (Pty) Ltd  
Co. Reg. No. 2002/021376/07  
P O Box 46, Isando 1600, RSA

**Marketed by:**

Afrivet Business Management (Pty) Ltd  
Co. Reg. No. 2000/011263/07  
P O Box 12406, Onderstepoort 0110, R.S.A.  
Tel: + 27 (0)12 545-0004