

FOR EXTERNAL ANIMAL USE ONLY

TRIATIX CATTLE POUR-ON

Reg. No. G3444 Act 36/1947

READY-TO- USE POUR-ON FOR THE CONTROL OF TICKS ON CATTLE

Contains: Amitraz 2,0 % m/v

WARNINGS

- Do not use on horses
- **DO NOT USE ON CALVES/LAMBS UNDER 1 MONTH.**
- **WITHDRAWAL PERIOD:**
 - Milk – 2 days
 - Meat – 7 days
- Signs of salivation and lethargy may occur with overdosage. These signs are transient and will pass with no ill effect.
- Keep out of reach of uninformed persons, children and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Do not eat, drink or smoke during or after handling this product, before washing hands and face thoroughly.
- Avoid contact with pour-on and wash immediately and thoroughly with soap and hot water after accidental contamination.
- Remove and wash all contaminated clothing.
- Destroy empty containers and do not re-use for any other purpose.
- Do not contaminate dams, rivers and other water sources.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

- **FOR EXTERNAL USE ON CATTLE ONLY**
- **Dose:** 1 ml/10 kg. Treat strictly according to body mass.
- TRIATIX CATTLE POUR-ON is a ready-to-use product and must not be mixed with any other dip, solvent or insecticidal compound.
- Apply in two lines along each side of the spine from the shoulder to the root of the tail.
- **In cases of heavy brown ear or blue tick challenge**, apply TRIATIX CATTLE POUR-ON in two lines from the base of the skull, along each side of the spine to the root of the tail.
- **In cases of heavy red-legged tick infestation**, it may be necessary to spot-treat the affected area under the tail above the anus.
- **Treat at regular intervals according to tick challenge.** The longer the product is regularly applied, the more the tick population is reduced. It will then become possible to apply at intervals greater than 7 days.

DOSE: Cattle - 1 ml/10 kg

200 ml	Treats 5 X 400 kg cattle
500 ml	Treats 12 X 400 kg cattle
1 l	Treats 20 X 400 kg cattle
5 l + 1 l	Treats 125 X 400 kg cattle
20 l	Treats 500 X 400 kg cattle

How to use the applicator

Remove the cap of the calibrated chamber. Without removing the remaining cap, squeeze the container until the correct dosage volume is present in the calibrated chamber, and apply as prescribed.

STORAGE INSTRUCTIONS

Protect from extremes of heat, cold and from dampness.

Store away from food and feed.

Tightly reseal the container if all the contents have not been used.

Presentation: 200ml, 500 ml and 1 l squeeze-pour bottles ; 5l and 20l refills

Registration holder:

Cooper Veterinary Products (Pty) Ltd

Co. Reg. No. 2002/021376/07

P O Box 46, Isando 1600, RSA

Marketed by:

Afrivet Business Management (Pty) Ltd

Co. Reg. No. 2000/011263/07

P O Box 12406, Onderstepoort 0110, R.S.A.

Tel: + 27 (0)12 545-0004