

FOR EXTERNAL ANIMAL USE ONLY

**TRIATIX 2 %
PIG POUR-ON**

Reg. No. G3381 Act 36/1947

Ready-to-use Emulsifiable Concentrate

STORAGE INSTRUCTIONS

Store below 25 °C in a dry place.

COMPOSITION

Amitraz 2 % m/v

INDICATIONS

Kills sarcoptic mange mites and lice on pigs.

WARNINGS

- DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 7 DAYS OF LAST APPLICATION.
- DO NOT USE ON HORSES.
- In case of poisoning call a doctor and make this information available.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions. Failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advise and notify the registration holder.

PRECAUTIONS

- Avoid contact with pour-on and wash immediately and thoroughly with soap and hot water after accidental contamination.
- Remove and wash contaminated clothing.
-
- Do not eat, drink or smoke during or after applying before washing hands and face thoroughly.
- Tightly reseal the container if all the contents have not been used.
- Destroy empty container by perforation and flattening and NEVER re-use for any other purpose.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

For external animal use only.

TRIATIX 2 % PIG POUR-ON is a ready-to-use product and must not be mixed with any other dip, solvent or insecticidal compound.

Apply to animals in a group whether visually infested or not.

Animals of all ages, including pregnant and suckling sows, suckers, piglets, weaners, growers, finishers and boars may be treated.

Apply the calculated amount of TRIATIX 2 % PIG POUR-ON evenly along the backline BETWEEN THE RUMP AND THE EARS.

In large boars and sows the total amount of TRIATIX 2 % PIG POUR-ON should be spread along the backline and sides to increase the spread of the remedy over the body.

The inside of the ears **MUST** be treated with a small amount of the calculated dose at each application. This is especially important in sows and boars.

Application rate

The dose will provide 5 – 6 mg of Amitraz per kg body mass and is calculated as quarter the body mass (kg) in ml e.g.

Body mass (kg)	Application (ml)
5	1,25
10	2,5
20	5,0
30	7,5
40	10,0
50	12,5
60	15,0
100	25,0

Initial application

Two applications to the entire herd with a 7 – 10 day interval.

In the case of severe infestation a third dose may be needed 10 days later.

Follow-up and subsequent applications

Apply to sows twice at weekly intervals within the last month before farrowing.

Apply to boars monthly using two applications at weekly intervals.

Recurrence in growers to be treated twice at weekly intervals.

All introductions to the unit to be treated three times at weekly intervals whilst in quarantine.

Culling of all badly infested sows and boars must be considered.

As TRIATIX 2 % PIG POUR-ON spreads over the surface of the pig's skin, killing parasites in the more inaccessible areas, there is no need to scrub the pig to remove scales before application.

Presentation: 1lt, 5lt

Registration holder:

Cooper Veterinary Products (Pty) Ltd

Co. Reg. No. 2002/021376/07

P O Box 46, Isando 1600, RSA

Marketed by:

Afrivet Business Management (Pty) Ltd

Co. Reg. No. 2000/011263/07

P O Box 12406, Onderstepoort 0110, R.S.A.

Tel: + 27 (0)12 545-0004