

FOR EXTERNAL ANIMAL USE ONLY

WIPE-OUT®
Reg. No. G3172 Act 36/1947
N-SR0858

READY-TO-USE, WATER-BASED POUR-ON FOR SHEEP AND GOATS

- Kills red lice.
- Controls Karoo paralysis ticks.
- Kills blowfly maggots.
- Kills keds on sheep.
- Contains no oil. Will not stain the fleece.

Contains: Deltamethrin 0,5 % m/v

WARNINGS

- DO NOT SLAUGHTER TREATED ANIMALS FOR HUMAN CONSUMPTION WITHIN 3 DAYS OF TREATMENT.
- DO NOT USE ON LACTATING SHEEP AND GOATS WHERE MILK IS INTENDED FOR HUMAN CONSUMPTION.
- DO NOT USE ON ANIMALS SEVERELY STRESSED OR IN ILL HEALTH,
- Toxic to fish and bees. DO NOT EMPTY INTO STREAMS.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Avoid contact with skin and eyes.
- Do not eat, drink or smoke while handling this product.
- Wash hands thoroughly after use.
- In case of accidental contact wash affected area with plenty of water and soap,

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

READY-TO-USE. SHAKE WELL.

Red lice on sheep and goats and keds on sheep

Sheep: 2 mℓ per 5 kg.

Goats: 2 treatments of 2 mℓ/5 kg applied 1 month apart provides 16 weeks (4 months) protection against red lice.

Simply apply the recommended dosage as set out below as a single strip along the backline using any dosing gun or applicator.

For best results use the Wipe-Out T-bar nozzle which enables rapid dispersion.

Control of red lice and keds is achieved in 3 - 4 weeks.

Wipe-Out can be applied to goats and British breeds of sheep with up to 6 months fleece growth or to Merinos with a maximum of 3 months of wool growth,

For optimum results treatment (April to June) just before the lice population peak (June to August) is recommended.

Treat all animals in the flock.

Dosage recommendations for red lice and keds

Body mass	Dose (mℓ)	Doses in 1 ℓ	Doses in 5 ℓ	Doses in 1 ℓ
Up to 10 kg	4	250	1 250	2 500
11 to 20 kg	8	125	625	1 250
21 to 30 kg	12	83	416	832
31 to 40 kg	16	62	312	624
41 to 50 kg	20	50	250	500
51 to 60 kg	24	41	208	416
61 to 70 kg	28	35	178	356
71 kg + over	30	33	166	332

Karoo paralysis ticks:

- **Woolled sheep:** 1 mℓ/5 kg. Repeat after 8 - 10 weeks if necessary.
- **Hairy sheep:** 2 mℓ/5 kg. Repeat after 4 weeks if necessary.
- **Goats:** 2 mℓ/5 kg. Repeat every 2 – 3 weeks if necessary.

Using any suitable applicator apply the recommended dosage rate given below divided into 4 equal parts to each axilla and groin e.g. a 60 kg sheep would receive a total of 12 mℓ given as 3 mℓ under each leg. For optimum results treatment just before the tick challenge is recommended.

Recommended dosage for Karoo paralysis tick control

Body mass	Total dosage (mℓ)			Divided dose per leg (mℓ)			Doses in 1 ℓ		
	Sheep		Goats	Sheep		Goats	Sheep		Goats
	Woolled	Hairy	-	Woolled	Hairy	-	Woolled	Hairy	-
Up to 10 kg	2	4	4	0,5	1,0	1,0	500	250	250
11 to 20 kg	4	8	8	1,0	2,0	2,0	250	125	125
21 to 30 kg	6	12	12	1,5	3,0	3,0	166	83	83
31 to 40 kg	8	16	16	2,0	4,0	4,0	125	62	62
41 to 50 kg	10	20	20	2,5	5,0	5,0	100	50	50
51 to 60 kg	12	24	24	3,0	6,0	6,0	83	41	41

61 to 70 kg	14	28	28	3,5	7,0	7,0	71	35	35
71 kg & over	16	32	32	4,0	8,0	8,0	62	31	31

Body mass	Total dosage (mℓ)			Divided dose per leg (mℓ)			Doses in 5 ℓ		
	Sheep		Goats	Sheep		Goats	Sheep		Goats
	Wooled	Hairy	-	Wooled	Hairy	-	Wooled	Hairy	-
Up to 10 kg	2	4	4	0,5	1,0	1,0	2 500	1 250	1 250
11 to 20 kg	4	8	8	1,0	2,0	2,0	1 250	625	625
21 to 30 kg	6	12	12	1,5	3,0	3,0	833	416	416
31 to 40 kg	8	16	16	2,0	4,0	4,0	625	312	312
41 to 50 kg	10	20	20	2,5	5,0	5,0	500	250	250
51 to 60 kg	12	24	24	3,0	6,0	6,0	416	208	208
61 to 70 kg	14	28	28	3,5	7,0	7,0	357	178	178
71 kg & over	16	32	32	4,0	8,0	8,0	312	156	156

Body mass	Total dosage (mℓ)			Divided dose per leg (mℓ)			Doses in 10 ℓ		
	Sheep		Goats	Sheep		Goats	Sheep		Goats
	Wooled	Hairy	-	Wooled	Hairy	-	Wooled	Hairy	-
Up to 10 kg	2	4	4	0,5	1,0	1,0	5 000	2 500	2 500
11 to 20 kg	4	8	8	1,0	2,0	2,0	2 500	1 250	1 250
21 to 30 kg	6	12	12	1,5	3,0	3,0	1 666	833	833
31 to 40 kg	8	16	16	2,0	4,0	4,0	1 250	625	624
41 to 50 kg	10	20	20	2,5	5,0	5,0	1 000	500	500
51 to 60 kg	12	24	24	3,0	6,0	6,0	832	416	416
61 to 70 kg	14	28	28	3,5	7,0	7,0	714	357	357
71 kg & over	16	32	32	4,0	8,0	8,0	624	312	312

- **Blowfly maggots:** Clip wool around the wound. Using any suitable applicator apply sufficient Wipe-Out to lightly cover the affected area and its immediate surroundings (± 10 mℓ). It will be necessary to monitor the area regularly and to retreat the area if necessary.

Presentation: 5 ℓ

Registration holder:

Cooper Veterinary Products (Pty) Ltd
Co. Reg. No. 2002/021376/07
P O Box 46, Isando 1600, RSA

Marketed by:

Afrivet Business Management (Pty) Ltd
Co. Reg. No. 2000/011263/07
P O Box 12406, Onderstepoort 0110, R.S.A.
Tel: + 27 (0)12 545-0004