

PACKAGE INSERT: AFRIVET REDLINE POUR-ON

FOR EXTERNAL ANIMAL USE ONLY

AFRIVET REDLINE POUR-ON

RSA Reg. No. G4245 Act 36/1947

Namibia  V04/18.3.3/538

HARMFUL

INDICATIONS

READY-TO- USE POUR-ON FOR TICKS ON CATTLE, SHEEP, OSTRICHES, AND GAME

- Controls ticks on cattle, sheep, ostriches, and game.
- Controls Tsetse fly worry on cattle.
- Controls feather mite on ostriches.
- Sterilises female ticks.
- Easy application.
- Residual action.
- Non-systemic action.
- The product is rainfast.

COMPOSITION

Flumethrin 1 % m/v

STORAGE INSTRUCTIONS

Protect from extremes of heat, cold and from dampness.

Store away from food and feed.

Tightly reseal the container if all the contents have not been used.

WARNINGS

- WITHDRAWAL PERIOD FOR MILK: NO WITHDRAWAL.
- WITHDRAWAL PERIOD FOR MEAT: 7 DAYS.
- Toxic to bees and fish.
- Although the colour of the product may vary, it will not influence the efficacy or residual period thereof.
- Skin reactions may occur in a small percentage of cases. If this happens, wash the affected animals with soap and water.
- Keep out of reach of uninformed persons, children and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Do not eat, drink or smoke during or after handling this product, before washing hands and face thoroughly.
- Avoid contact with pour-on and wash immediately and thoroughly with soap and hot water after accidental contamination.
- Remove and wash all contaminated clothing.
- Destroy empty containers and do not re-use for any other purpose.

SLEGS VIR UITWENDIGE DIERGEBRUIK

AFRIVET REDLINE OPGIETMIDDEL

RSA Reg. No. G4245 Act 36/1947

Namibia  V04/18.3.3/538

SKADELIK

AANWYSINGS

GEREED-VIR-GEBRUIK OPGIETMIDDEL VIR BOSLUISE OP BEESTE, SKAPE, VOLSTRUISE EN WILD

- Beheer bosluise op beeste, skape, volstruise en wild.
- Beheer Tsetsevliegplae by beeste.
- Beheer veermyte op volstruise.
- Steriliseer wyfiebosluise.
- Maklike toediening.
- Nawerking.
- Nie-sistemiese aksie.
- Die produk is reënvas.

SAMESTELLING

Flumetriën 1 % m/v

BERGINGSAAANWYSINGS

Beskerm teen uiterstes van hitte, koue en teen klammigheid.

Berg weg van voedsel en voer.

Indien inhoud nie opgebruik word nie, herseël houër deeglik.

WAARSKUWINGS

- ONTTREKKINGSPERIODE VIR MELK: GEEN.
- ONTTREKKINGSPERIODE VIR VLEIS: 7 DAE.
- Giftig vir bye en visse.
- Alhoewel die produk se kleur mag variëer, sal dit geen invloed hê op die doeltreffendheid of nawerking van die produk nie.
- Velreaksie mag in 'n klein persentasie van gevalle voorkom. Indien dit gebeur, was die geaffekteerde diere met seep en water.
- Hou buite bereik van oningeligte persone, kinders en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

VOORSORGMATREËLS

- Moenie eet, drink of rook tydens of na aanwending van die middel alvorens die hande en gesig deeglik gewas is nie.
- Vermoedlik aanraking met opgietsmiddel en was onmiddellik en deeglik met seep en warm water na toevallige besmetting.
- Verwyder en was besoedelde klere.
- Vernietig leë houers en moenie vir enige ander doel gebruik nie.
- Moenie damme, riviere en ander waterbronne besoedel nie.

- Do not contaminate dams, rivers and other water sources.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

- Treat at regular interval according to tick challenge.
- The longer the product is regularly applied, the more the tick population is reduced. It may then become possible to treat at extended intervals.

Dosage

The basic dosage is 1 ml per 10 kg body mass (refer to dosage table below).

- **Cattle:** Apply the required dose in two lines, from the base of the skull along both sides of the spine up to the tail root. For Tsetse fly treat every 14 days or according to challenge.
- **Sheep:** Apply the required dose volume in 4 parts to the bare skin areas of the groin and axillae. For **prevention** of Karoo Paralysis, Afrivet Redline has to be applied early in the season when the Karoo Paralysis tick is first observed (either on the grazing or on the animals). Two to three treatments 7 to 10 days apart at this time should protect the sheep against Karoo Paralysis, depending on the severity of the tick challenge. For **treatment** of Karoo Paralysis (when the sheep are already recumbent), pour-on dip formulations are not recommended, as it takes a while for the product to distribute over the skin. Ensure immediate detachment of **all** ticks by spraying the belly and legs of affected animals with an amitraz-based product such as Eraditick 125.
- **Game:** Apply the required dose volume with a suitable applicator. If self-medicating devices are used, it is important that the device only remains activated for as long as it takes to deliver one dosage to the animals in a particular paddock / camp. The device should not be used for continuous, low-level treatment.
- **Ostriches:** Apply the required dose volume along the backline.

**Dosage table: Cattle and Game/
Dosistabel: Beeste en Wild**

Body mass (kg)	Dosage	No. of doses per 1 ℓ
Up to 100 kg	1 ml per 10 kg	
100 kg	10 ml	100
200 kg	20 ml	50
300 kg	30 ml	33
400 kg	40 ml	25
500 kg and over	50 ml	20

Applicator: Squeeze bottle with dose meter. Squeeze required volume into dose meter or use a drench gun applicator.

Aanwender: Druk bottel met dosismeter totdat vereiste volume in die dosismeter is, of gebruik 'n aanwender.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

- Behandel met gereelde tussenposes na gelang van die bosluisuitdaging.
- Hoe langer die produk gereeld gebruik word, hoe groter die vermindering in die bosluisbevolking. Dit mag dan moontlik word verlengde periodes tussen behandelinge te hê.

Dosis

Die basiese dosis is 1 ml/10 kg liggaamsmassa (verwys na doseringstabel onder).

- **Beeste:** Dien die vereiste dosis toe in twee strepe, vanaf die basis van die kop langs beide kante van die ruggraat tot op die stertwortel. Vir Tsetsevlieg, behandel met 14 dae tussenposes of na gelang van die vliegplaag.
- **Skape:** Dien die vereiste dosis in 4 dele toe aan die gedeeltes van die lieste en oksels waar die vel kaal is. Om Karooverlamming te **voorkom**, moet Afrivet Redline vroeg in die seisoen aangewend word wanneer die bosluise vir die eerste maal opgemerk word (op die weiding of op die diere). Twee of drie behandelings met 'n 7 tot 10 dae tussenpose op hierdie tyd behoort voldoende beskerming teen Karooverlamming te bied afhangend van die graad van bosluisbesmetting. Om Karooverlamming te **behandel** (wanneer die skape alreeds lê), word opgietsformulasies nie aanbeveel nie omrede die produk 'n tyd lank neem om oor die vel te versprei. Verseker onmiddellike loslating van **alle** bosluise deur die pense en bene van besmette diere te bespuit met 'n amitraz-gebaseerde produk soos Eraditick 125.
- **Wild:** Dien die vereiste dosis met 'n gepaste toediener toe. Indien 'n selftoediener gebruik word, is dit belangrik dat die toestel slegs geaktiveer bly vir solank dit nodig is om die diere in 'n betrokke kamp eenmaal te behandel. Die toestel moet nie vir langdurige, laevlak behandeling gebruik word nie.
- **Volstruise:** Wend die vereiste dosis met die ruglyn langs aan.

**Dosage table: Sheep and Ostriches/
Dosistabel: Skape en Volstruise**

Body mass (kg)	Dosage	No. of doses per 1 ℓ
20	2 ml	500
30	3 ml	333
40	4 ml	250
50	5 ml	200
60	6 ml	166

PRESENTATION/AANBIEDING

10 ml, 20 ml, 40 ml, 200 ml, 500 ml, 1 L, 5 L, 20 L

Registration holder/Registrasiehouer:

Afrivet Business Management (Pty) Ltd

Co./Mpy. Reg. No. 2000/011263/07

P O Box 2009, Faerie Glen

0043, Pretoria

Tel: +27 12 817 9060

Helpline: 0860 833 2787