

PACKAGE INSERT: ECOMINTIC 50

FOR ANIMAL USE ONLY

ECOMINTIC 50

RSA Reg. No. G2000 Act 36/1947

Namibia S0 V02/18.1.1/1146

②†

CAUTION

STORAGE INSTRUCTIONS

Store in a cool, dry place.

COMPOSITION

Fenbendazole 5 % m/v.

INDICATIONS

A remedy against roundworms in cattle, horses and ostriches and roundworms and milk tapeworm in sheep and goats.

WARNINGS

- DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 7 DAYS OF TREATMENT.
- Do not mix or dilute with any other substance.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large range of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- It is recommended that sheep be vaccinated against pulpy kidney disease before any worm remedy is administered.

DIRECTIONS FOR USE – USE ONLY AS DIRECTED

Use as an oral drench.

Shake well before use.

Dosage rate:

Cattle: 1 ml/10 kg body mass.

Horses: 2 ml/10 kg body mass.

Sheep and Goats:

- **Roundworms** 1 ml/10 kg body mass.

- **Milk tapeworm (*M. expansa*)** 2 ml/10 kg body mass.

Ostriches: 3 ml/10 kg body mass.

Example for cattle and horses (roundworms):

Body mass (kg)	Dose (ml)		No. of doses in pack			
	Cattle	Horses	Cattle		Horses	
			200 ml	500 ml	200 ml	500 ml
50	5	10	40	100	20	50
100	10	20	20	50	10	25
150	15	30	13	33	6	16
200	20	40	10	25	5	12
250	25	50	8	20	4	10
300	30	60	6	16	3	8
350	35	70	5	14	2	7

Example for sheep, goats and ostriches (roundworms and milk tapeworm):

Body mass (kg)	Dose (ml)			No. of doses in pack					
	Sheep and Goats		Ostriches	Sheep and Goats				Ostriches	
	Roundworm	Milk Tapeworm (<i>M. expansa</i>)		Roundworm		Milk Tapeworm (<i>M. expansa</i>)			
				200 ml	500 ml	200 ml	500 ml	200 ml	500 ml
10	1	2	3	200	500	100	250	66	166
20	2	4	6	100	250	50	125	33	83
30	3	6	9	66	166	33	83	22	55
40	4	8	12	50	125	25	62	16	41
50	5	10	15	40	100	20	50	13	33

EFFICACY

• **Cattle, sheep and goats**

Worm species	Immature	Adult
• Cattle		
Wireworm	*	*
Brown stomachworm	*	*
Nodular worm	*	*
Hookworm	*	*
Lungworm	**	*
• Sheep and Goats		
Wireworm	*	*
Brown stomachworm	*	*
Bankruptworm	*	*
Long-necked bankruptworm	*	*
White bankruptworm	*	*
Hookworm	*	*
Large-mouthed bowelworm	*	*
Nodular worm	*	*
Lungworm	*	*
Milk tapeworm (<i>Moniezia expansa</i>)	Class I	

* Control = ≥ 90 % effective

** Aids in Control = 60 – 89 % effective

Class I: 100 % effective in more than 80 % of treated animals.

Kills worm eggs present in the animal at the time of dosing.

- **Horses**

- * Bloodworms (large strongyles), redworms (small strongyles), pinworm (*Oxyuris*), ascarids and stomach worm (*Habronema majus*).

- **Ostriches**

- * *Libyostrongylus douglassii*
- * Control = ≥ 90 % effective

The list contains the more important endoparasite species. This anthelmintic may also be effective against other species. For more information consult your veterinarian.

† Benzimidazole

Packaging Size:

200 ml, 500 ml, 1 l, 2 l, 5 l, 10 l and 20 l

Registration holder:

ECO Animal Health Southern Africa (Pty) Ltd
Co. Reg. No. 1992/000835/07
P O Box 2009, Faerie Glen
0043, R.S.A.
Helpline: 0860 833 2787
www.ecoanimalhealth.com

Distributed by:

Afrivet Business Management (Pty) Ltd
Co. Reg 2000/011263/07
PO Box 2009, Faerie Glen
0043, RSA
Helpline: 0860 833 2787
www.afrivet.co.za

VOUBILJET: ECOMINTIC 50

SLEGS VIR DIEREGERBRUIK

ECOMINTIC 50

RSA Reg. No. G2000 Wet 36/1947

Namibia S0 V02/18.1.1/1146

②†

VERSIGTIG

BERGINGSANWYSINGS

Bewaar in 'n koel, droë plek.

SAMESTELLING

Fenbendasool 5 % m/v.

AANDUIDINGS

'n Middel teen rondewurms in beeste, perde en volstruise en rondewurms en melklintwurm in skape en bokke.

WAARSKUWINGS

- MOENIE DIERE VIR MENSLIKE VERBRUIK BINNE 7 DAE NA BEHANDELING SLAG NIE.
- Moenie met enige ander middel vermeng nie.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

VOORSORGMAATREËLS

- Dit word aanbeveel dat alle skape teen bloednier ingeënt word voordat dosering met enige wurmmiddel plaasvind.

GEBRUIKSAANWYSINGS – GEBRUIK SLEGS SOOS AANGEDUI

Gebruik as 'n mondelingse dosering.

Skud deeglik voor gebruik.

Dosis:

Beeste: 1 ml/10 kg liggaamsmassa.

Perde: 2 ml/10 kg liggaamsmassa.

Skape en Bokke:

• **Rondewurms** 1 ml/10 kg liggaamsmassa.

• **Melklintwurm (*M. expansa*)** 2 ml/10 kg liggaamsmassa.

Volstruise: 3 ml/10 kg liggaamsmassa.

Voorbeeld vir beeste en perde (rondewurms):

Liggaamsmassa (kg)	Dosis (ml)		Aantal dosisse in verpakking			
	Beeste	Perde	Beeste		Perde	
			200 ml	500 ml	200 ml	500 ml
50	5	10	40	100	20	50
100	10	20	20	50	10	25
150	15	30	13	33	6	16
200	20	40	10	25	5	12
250	25	50	8	20	4	10
300	30	60	6	16	3	8
350	35	70	5	14	2	7

Voorbeeld vir skape, bokke en volstruise (rondewurms en melklintwurm):

Liggaamsmassa (kg)	Dosis (ml)			Aantal dosisse in verpakking					
	Skape en Bokke		Volstruise	Skape en Bokke				Volstruise	
	Rondewurms	Melklintwurm (<i>M. expansa</i>)		Rondewurms		Melklintwurm (<i>M. expansa</i>)			
				200 ml	500 ml	200 ml	500 ml	200 ml	500 ml
10	1	2	3	200	500	100	250	66	166
20	2	4	6	100	250	50	125	33	83
30	3	6	9	66	166	33	83	22	55
40	4	8	12	50	125	25	62	16	41
50	5	10	15	40	100	20	50	13	33

DOELTREFFENDHEID

• **Beeste, skape en bokke**

Wurmsoort	Onvolwasse	Volwasse
• Beeste		
Haarwurm	*	*
Bruinmaagwurm	*	*
Knoppieswurm	*	*
Haakwurm	*	*
Longwurm	**	*
• Skape en bokke		
Haarwurm	*	*
Bruinmaagwurm	*	*
Bankrotwurm	*	*
Langnekbankrotwurm	*	*
Witbankrotwurm	*	*
Haakwurm	*	*
Grootbekwurm	*	*
Knoppieswurm	*	*
Longwurm	*	*
Melklintwurm (<i>Moniezia expansa</i>)	Klas I	

* Beheer = ≥ 90 % effektief

** Hulp in beheer = 60 – 89 % effektief

Klas I: 100 % doeltreffend in meer as 80 % van behandelde diere.

Dood wurmeiers wat tydens dosering in die dier teenwoordig is.

- **Perde**

- * Bloedwurms (groot strongiele), rooiwurms (klein strongiele), sambokwurm (*Oxyuris*), spoelwurm (askariede) en maagwurm (*Habronema majus*).

- **Volstruise**

- * *Libyostrongylus douglassii*.

* Beheer = ≥ 90 % effektief.

Die lys bevat die meer belangrike endoparasitiese spesies. Dit kan ook effektief teen ander spesies gebruik word. Vir verdere inligting raadpleeg u veearts.

† Bensimidiasool

AANBIEDING:

200 ml, 500 ml, 1 l, 2 l, 5 l, 10 l and 20 l

Registrasiehouer:

ECO Animal Health Southern Africa (Pty) Ltd
Mpy. Reg. No. 1992/000835/07
Posbus 2009, Faerie Glen
0043, R.S.A.
Hulplyn: 0860 833 2787

Versprei deur:

Afrivet Business Management (Pty) Ltd
Mpy. Reg 2000/011263/07
Posbus 2009, Faerie Glen
0043, RSA
Hulplyn: 0860 833 2787