

PROPOSED LABEL: ERADITICK® PLUS POUR-ON

FOR EXTERNAL ANIMAL USE ONLY

ERADITICK® PLUS POUR-ON

Reg. No. G4251 Act 36/1947

Namibia [SQ] V04/18.3.9/541

READY-TO-USE POUR-ON FOR CATTLE, SHEEP AND GOATS.

- RESIDUAL ACTION.
- NON-SYSTEMIC ACTION.

Cattle

- EASY APPLICATION ALONG THE BACKLINE.
- Controls ticks.
- Controls stable flies, horn flies, cattle louse flies and nuisance flies e.g. house flies.
- Kills lice (biting and sucking) and mange mites.
- Protects against black flies.

Sheep and goats

- Controls Karoo-paralysis, bont-legged and red-legged ticks.

CAUTION

COMPOSITION

Amitraz	1,5 % m/v
Deltamethrin	0,50 % m/v
Piperonyl butoxide	3,0 % m/v

STORAGE INSTRUCTIONS

- Store in a cool place.
- Store away from food and feed.
- Tightly reseal the container if all the contents have not been used.

WARNINGS

- Withdrawal periods:
 - Meat: 7 days.
 - Milk: 0 days.
- DO NOT USE ON HORSES.
- DO NOT USE ON CALVES/LAMBS UNDER 3 MONTHS OF AGE.
- Treat strictly according to body mass.
- Signs of salivation and lethargy may occur with overdosing. These signs are transient and will pass with no ill effect.
- Pyrethroid pour-ons may cause irritation especially in dairy cattle. It is suggested that a group of 10 animals be treated first and observed for an hour before treating the rest of the herd.
- Keep out of reach of children, uninformed persons and animals.

SLEGS VIR UITWENDIGE DIEREGERBUK

ERADITICK® PLUS OPGIETMIDDEL

Reg. No. G4251 Wet 36/1947

Namibia [SQ] V04/18.3.9/541

GEREED-VIR-GERBUK OPGIETMIDDEL VIR BEESTE, SKAPE EN BOKKE.

- NAWERKENDE EFFEK.
- 'N NIE-SISTEMISE WERKING.

Beeste

- MAKLIKE TOEDIENING LANGS DIE RUGLYN.
- Beheer bosluise.
- Beheer stalvlieë, horing-, beesluis- en lasvlieë bv. huisvlieë.
- Dood luise (bytende en suigende) en skurftemyte.
- Beskerm teen riviermuggies.

Skape en bokke

- Beheer Karooverlamming-, bontpoot- en rooipootbosluise.

VERSIGTIG

SAMESTELLING

Amitraz	1,5 % m/v
Deltamethrin	0,50 % m/v
Piperonylbutoksied	3,0 % m/v

BERGINGSAAWYSINGS

- Berg in 'n koel plek.
- Berg weg van voedsel en voer.
- Indien inhoud nie opgebruik word nie, herseël houer deeglik.

WAARSKUWINGS

- Onttrekkingsperiodes:
 - Vleis: 7 dae.
 - Melk: 0 dae.
- MOENIE OP PERDE GERBUK NIE.
- MOENIE OP KALWERS/LAMMERS JONGER AS 3 MAANDE GERBUK NIE.
- Behandel streng volgens liggaamsmassa.
- Tekens van kwyling en lomerigheid mag voorkom met oordosering. Hierdie tekens is tydelik en diere sal herstel sonder enige nuwe-effek.
- Piretroïdeopgiemiddels mag irritasie veroorsaak, veral in melkbeeste. Dit word aanbeveel om 'n groep van 10 diere te behandel en hulle vir 'n uur waar te neem voordat die res van die kudde behandel word.
- Hou buite die bereik van kinders, oningeligte persone en diere.

- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Do not eat, drink or smoke during or after handling this product, before washing hands and face thoroughly.
- Avoid contact with pour-on and wash immediately and thoroughly with soap and hot water after accidental contamination.
- Remove and wash all contaminated clothing.
- Destroy empty containers and do not re-use for any other purpose.
- Do not contaminate dams, rivers and other water sources.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

- SHAKE WELL BEFORE USE.
- ERADITICK® PLUS POUR-ON is a ready-to-use product and must not be mixed with any other dip, solvent or insecticidal compound.

Cattle:

- **Ticks, flies and lice:** Dose 1 ml / 10 kg.
- Apply in two lines, from the base of the skull down both sides of the spine up to the tail root. Application at the base of the skull is especially important in the case of heavy Brown ear-tick infestation.
- Treat at regular intervals according to tick challenge. The longer the product is applied at regular intervals, the more the tick population is reduced. It will then become possible to apply at intervals greater than 7 days.
- **For Black fly (*Simulium spp*)** apply also to the ears and underline. Repeat as required according to challenge.
- **For control of nuisance flies and stable flies**, use 2 – 3 treatments at weekly intervals. Thereafter use according to challenge.

Sheep and goats

- Environmental conditions and parasite challenge can influence treatment intervals.
- **Dosage for optimal residual action:**
1 ml / 5 kg
- **Karoo paralysis ticks:**
Divide the full dose per animal into 4 equal parts. Apply one part onto each axillae and groin region, whilst the animal is turned over in a sitting position.
Frequency:
Angora goats – repeat after ± 2 weeks.
Dorper sheep – repeat after ± 4 weeks.
Merino sheep – repeat after ± 6 weeks.
- **Bont-legged and red-legged ticks:**

- Alhoewel hierdie produk breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

VOORSORGMATREËLS

- Moenie eet, drink of rook tydens of na aanwending van die middel alvorens die hande en gesig deeglik gewas is nie.
- Vermoedlik aanraking met opgietsmiddel en was onmiddellik en deeglik met seep en warm water na toevallige besmetting.
- Verwyder en was besoedelde klere.
- Vernietig leë houers en moenie vir enige ander doel gebruik nie.
- Moenie damme, riviere en ander waterbronne besoedel nie.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

- SKUD DEEGLIK VOOR GEBRUIK.
- ERADITICK® PLUS OPGIETMIDDEL is 'n gereed-om-te-gebruik produk en moet met geen ander dip, oplossing of insekdodende middel gemeng word nie.

Beeste:

- Bosluise, vlieë en luise: Doseer 1 ml / 10 kg.
- Dien die nodige dosis toe in twee strepe, vanaf die basis van die skedel langs beide kante van die ruggraat tot by die stertwortel. Toediening by die basis van die skedel is veral belangrik met erge Bruinoorbosluislus.
- Behandel met gereelde tussenposes, na gelang van bosluispopulasie. Hoe langer die produk gereeld gebruik word, hoe groter die daling in die bosluisbevolking. Dit mag dan moontlik word om die tussenpose van 7 dae tussen behandelings te verleng.
- Vir riviermuggies (*Simulium spp*) wend ook aan ore en onderlyn. Herhaal soos benodig na gelang van die besmetting.
- Vir populasiebeheer van las- en stalvlieë gebruik 2– 3 behandelings met weeklikse tussenposes. Gebruik daarna na gelang van besmetting.

Skape en bokke

- Omgewingsfaktore en parasietuitdaging kan behandelingstussenposes beïnvloed.
- **Dosis vir optimale nawerkende effek:**
1 ml / 5 kg
- **Karoooverlammingsbosluise:**
Verdeel die volle dosis per dier in 4 gelyke dele. Wend een deel aan elke oksel en lies gedeelte, terwyl die dier omgekeer is in 'n sittende posisie.
Tussenpose:
Angorabokke – herhaal na ± 2 weke.
Dorperskape – herhaal na ± 4 weke.
Merinoskape – herhaal na ± 6 weke.

The dosage should be divided into 3 equal parts and applied to the axillae, groin and under the tail regions.

- Treat sheep at regular intervals of \pm 3 – 4 weeks or as necessary, according to tick challenge.
- For crippled sheep apply 1 ml to the affected hoof/fetlock.

Use of applicator

Remove the cap of the calibrated chamber. Without removing the remaining cap, squeeze the container until the correct dosage volume is present in the calibrated chamber, and apply as prescribed.

PRESENTATION

200 ml, 500 ml, 1 L, 5 L, 20 L

REGISTRATION HOLDER:

Afrivet Business Management (Pty) Ltd
Co. Reg. No. 2000/011263/07
P O Box 2009, Faerie Glen
0043, Pretoria
Tel: +27 12 817 9060
Helpline: 0860 933 2787
www.afrivet.co.za

- **Bontpoot- en rooipootbosluise:**

Verdeel die dosis in 3 gelyke dele en wend aan die oksel-, lies- en onder die stertgedeeltes.

- Behandel skape met gereelde tussenpose van ongeveer 3 – 4 weke of wanneer benodig, na gelang van bosluispopulasie.
- Vir skape wat kruppel is, wend 1 ml op die aangetaste hoef/koot aan.

Gebruik van toediener

Verwyder die prop van die gekalibreerde maatkamer. Sonder om die ander prop te verwyder, druk die toediener totdat die verlangde dosisvolume in die maatkamer versamel en dien toe soos voorgeskryf.

AANBIEDING

200 ml, 500 ml, 1 L, 5 L, 20 L

REGISTRASIEHOUER:

Afrivet Business Management (Pty) Ltd
Mpy. Reg. No. 2000/011263/07
Posbus 2009, Faerie Glen
0043, Pretoria
Tel: +27 12 817 9060
Hulplyn: 0860 933 2787
www.afrivet.co.za