

Package Insert: EXPEL™

EXPEL™

RSA Reg. No. G3245 Act 36/1947

Namibia  V04.18.3.3/482

Zambia: 335/707V

Botswana: W130581

STORAGE INSTRUCTIONS

STORE BETWEEN 19 – 25 °C.

COMPOSITION

Deltamethrin 0,10% m/m

INDICATIONS

A multi-purpose remedy which effectively controls and prevents screw-worm/blowfly strike, promotes healing and kills ticks present at time of application.

For use on livestock, horses, wild life, dogs, poultry and birds.

WARNINGS

- DO NOT APPLY TO CATS.
- APPLY SPARINGLY TO BIRDS.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Avoid inhalation of aerosol spray.
- Wash contaminated skin with soap and water.
- In the case of eye contact rinse immediately with copious amounts of water.
- In case of poisoning call a doctor and make this information available.
- Do not contaminate water sources or the environment.
- Dispose of containers in a safe manner and do not re-use for any other purpose.
- Store away from food and feed.

DIRECTIONS FOR USE – USE ONLY AS DIRECTED
DO NOT APPLY TO DEEP PENETRATING WOUNDS.

• **All wounds**

Apply in, around and above wounds (to diffuse downwards), to prevent screw-worm/blowfly strike and reduce bacterial infection and promote healing. This is recommended following routine procedures such as dehorning, knife castration, Mules operation and tail docking.

• **Screw-worm/blowfly strike treatment**

Apply freely into, around and above infested wounds. The worms will emerge within a short period, fall to the ground and die.

With small wounds one treatment ± every 4 days until healed.

Deeper wounds with extensive damage will respond more quickly if crusts and debris are gently removed before

EXPEL™

RSA Reg. No. G3245 Wet 36/1947

Namibia  V04.18.3.3/482

Zambia: 335/707V

Botswana: W130581

BERGINGSAAWYSINGS

BERG TUSSEN 19 – 25 °C.

SAAMSTELLING

Deltametrien 0,10% m/m

AANDUIDINGS

’n Veeldoelige middel vir doeltreffende beheer en voorkoming van spykerwurm/brommer-maaier aanvalle, help met genesing en dood bosluise teenwoordig tydens aanwending.

Vir gebruik op vee, perde, wild, honde, pluimvee en voëls.

WAARSKUWINGS

- MOENIE NIE OP KATTE GEBRUIK NIE.
- GEBRUIK SPAARSAAM OP VOËLS.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie middel breedvoerig onder ’n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg ’n veearts en verwittig die registrasiehouer.

VOORSORGMAATREËLS

- Vermyn inaseming van aërosol sproei.
- Was besoedelde kleres met seep en water.
- In geval van oogkontak spoel onmiddellik met baie water.
- In geval van vergiftiging raadpleeg ’n dokter en maak hierdie inligting beskikbaar.
- Moenie waterbronne of die omgewing besoedel nie.
- Doen weg met houers op ’n veilige manier en moenie vir enige ander doel hergebruik nie.
- Berg weg van kos en voer.

GEBRUIKSAANWYSINGS – GEBRUIK SLEGS SOOS AANGEDUI

MOENIE AAN DIEP DEURDRINGENDE WONDE AANWEND NIE.

• **Alle wonde**

Wend aan in, om en bokant wonde (vir afwaartse verspreiding), vir voorkoming van spykerwurm/brommer-maaier aanvalle, vermindering van bakteriese infeksie en help met genesing.

Bogenoemde word aanbeveel na prosedures soos onthoring, kastrasie, Mules operasie en stert afsit.

• **Behandeling vir spykerwurm/brommer-maaier aanvalle**

Gebruik ruimskoots in, om en bokant besmette wonde. Die wurms sal na ’n kort ruk verskyn, afval en vrek.

Vir klein wonde, een behandeling ± elke 4 dae totdat dit genees is.

Package Insert: EXPEL™

application but this is not essential.
Better results follow application to animals which are dry.

PRESENTATION:

300 ml

REGISTRATION HOLDER:

Afrivet Business Management (Pty) Limited
Co.Reg. No. 2000/011263/07
P O Box 2009
FAERIE GLEN, 0043 R.S.A.
Tel: 012 991 6416
Helpline: 0860 833 2787
Website: www.afrivet.co.za

Dieper wonde met baie beskadiging sal beter reageer as rowe en vuilgoed sagkens verwyder word voor aanwending, maar dit is nie noodsaaklik nie.

Beter resultate sal verkry word as die middel op droë diere aangewend word.

AANBIEDING

300ml

REGISTRASIEHOUER:

Afrivet Business Management (Pty) Limited
Mpy. Reg. No. 2000/011263/07
Posbus 2009
FAERIE GLEN, 0043 R.S.A.
Tel: 012 991 6416
Hulplyn: 0860 833 2787
Webblad: www.afrivet.co.za