

Package Insert: Moxziben Sheep Drench

FOR ANIMAL USE ONLY

MOXZIBEN SHEEP DRENCH ② + ⑧ + ① †
Reg. No. G4040 Act 36/1947

CAUTION

STORAGE INSTRUCTIONS

- Store in a cool, dry, well ventilated place away from sources of ignition and heat.
- Store only in original container and keep tightly closed.
- Keep away from food, drink and animal feed.

INDICATIONS

Deworming drench for sheep for the control of roundworms i.e. Wireworm *, Liver fluke, Brown stomach worm*, Bankruptworm, Long-necked bankruptworm and Large-mouthed bowelworm as well as Milk tapeworm. Controls nasal bot larvae infestation.

* **Persistency: Wireworm and Brown stomach worm 28 days.**

COMPOSITION

Triclabendazole	5.00 % m/v
Praziquantel	3.75 % m/v
Moxidectin	0.10 % m/v

WARNINGS

- **DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 10 DAYS OF LAST TREATMENT.**
- **DO NOT USE IN SHEEP PRODUCING MILK FOR HUMAN CONSUMPTION.**
- Do not administer to lambs under the age of 4 months or emaciated sheep.
- Do not administer to anaemic sheep. If the animals are anaemic, treat first with an anthelmintic product from another group. Consult a veterinarian for further information.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

SLEGS VIR DIEREGBRUIK

MOXZIBEN SHEEP DRENCH ② + ⑧ + ① †
Reg. No. G4040 Wet 36/1947

VERSIGTIG

BERGINGSAAWYSINGS

- Berg in 'n koel, droë, goed geventileerde plek. Vermy bronne van ontbranding en hitte.
- Berg alleenlik in oorspronklike houers en hou dit toe.
- Bêre weg van kos, drink en diere voedsel.

AANWYSINGS

Ontwurmingsmiddel vir skape vir die beheer van rondewurms d.w.s. Haarwurm*, Lewerslak, Bruinmaagwurm*, Bankrotwurm, Langnek-bankrotwurm e Grootbekwurm sowel as Melk-lintwurm. Beheer neuswurm larwe infestasië.

* **Nawerking: Haarwurm en Bruinmaagwurm 28 dae.**

SAMESTELLING

Triclabendazole/Triklabendazool	5.00 % m/v
Praziquantel/Prasikwantel	3.75 % m/v
Moxidectin/Moksidektien	0.10 % m/v

WAARSKUWINGS

- **MOENIE DIERE VIR MENSLIKE VERBRUIK BINNE 10 DAE NA LAASTE BEHANDELING SLAG NIE.**
- **MOENIE IN SKAPE GEBRUIK WAT MELK VIR MENSLIKE VERBRUIK PRODUSEER NIE.**
- Moenie lammers onder die ouderom van 4 maande of uitgeteerde skape doseer nie.
- Moet nie vir skape met bloedarmoede toedien nie. Indien diere bloedarmoedig is, behandel eers met 'n middel wat aan 'n verskillende wurmmiddel groep behoort. Raadpleeg u veearts vir meer inligting
- Hou buite die bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie produk breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

Package Insert: Moxziben Sheep Drench

PRECAUTIONS

- Sheep should be vaccinated against pulpy kidney (enterotoxaemia) at least 3 weeks prior to treatment. Consult your veterinarian.
- Avoid contact with skin and eyes.
- Wear personal protective equipment i.e. gloves, clothing, eye and face protection.
- Do not breathe fumes.
- Do not eat, drink or smoke while handling the product.
- Wash hands thoroughly after use.
- In case of ingestion consult a doctor and make this label available.
- Do not discharge into drains or the environment. If the product has contaminated water, inform the appropriate authorities. Moxidectin is harmful to fish and certain aquatic organisms.
- Dispose of left over product and its container at a hazardous- or special waste collection point. Do not re-use container for any other purpose.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

- SHAKE WELL BEFORE USE.
- MOXZIBEN SHEEP DRENCH is a ready-to-use suspension. Do not dilute or mix with any other product.

Dose

Roundworms, Liver fluke, Milk tapeworm and nasal bot: 1 ml/5 kg e.g.	
Body mass (kg)	Dose (mℓ)
Up to 20 kg	4.0
30	6.0
40	8.0
50	10.0
60	12.0

VOORSORGMATREËLS

- Skape behoort ten minste 3 weke voor behandeling teen bloednier (enterotoksemie) ingeënt te word. Raadpleeg 'n veearts.
- Vermoed kontak met vel en oë.
- Dra persoonlike beskermende toerusting d.w.s. handskoene, klere, oog en gesig beskerming.
- Moenie damp inasem nie.
- Moenie eet, drink of rook tydens hantering van die produk nie.
- Was hande deeglik na gebruik.
- In geval van inname raadpleeg 'n dokter en maak hierdie etiket beskikbaar.
- Moenie in riool of die omgewing weggooi nie. As die produk water besoedel het meld aan by die betrokke owerheid. Moxidectin is skadelik vir vis en sekere waterorganismes.
- Verwyder oorblywende produk en die houër by 'n skadelike- of spesiale afval verwyderings punt. Moenie houër vir enige ander doel hergebruik nie.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

- SKUD GOED VOOR GEBRUIK
- MOXZIBEN SHEEP DRENCH is 'n gereed- vir- gebruik suspensie. Moenie verdun of met ander produkte meng nie.

Dosis

Rondewurms, Lewerslak, Melkintwurm en neuswurm: 1 ml/5 kg b.v	
Liggaamsamassa (kg)	Dosis (mℓ)
Tot 20 kg	4.0
30	6.0
40	8.0
50	10.0
60	12.0

EFFICACY Roundworms	Adult/ Volwasse	DOELTREFFENDHEID Rondewurms
Wireworm (<i>Haemonchus contortus</i>) Persistency 28 days.	*	Haarwurm (<i>Haemonchus contortus</i>) Nawerking 28 dae.
Brown stomach worm (<i>Teladorsagia circumcincta</i>) Persistency 28 days.	*	Bruinmaagwurm (<i>Teladorsagia circumcincta</i>) Nawerking 28 dae.
Bankruptworm (<i>Trichostrongylus colubriformis</i>)	*	Bankrotwurm (<i>Trichostrongylus colubriformis</i>)
Long-necked bankruptworm (<i>Nematodirus spathiger</i>)	*	Langnekbankrotwurm (<i>Nematodirus spathiger</i>)
Large-mouthed bowelworm (<i>Chabertia ovina</i>)	*	Grootbekwurm (<i>Chabertia ovina</i>)

* Controls/Beheer = > 90% effective/doeltreffend

Package Insert: Moxziben Sheep Drench

EFFICACY Liver fluke					DOELTREFFENDHEID Lewerslak
Age	2	4	6	Adult/ Volwasse	Ouderdom
Liver Fluke (<i>F hepatica</i>)	*	*	*	*	Lewerslak (<i>F hepatica</i>)
	3	6	8	Adult/ Volwasse	
Giant Liver Fluke (<i>F gigantica</i>)	*	*	*	*	Reuse Lewerslak (<i>F gigantica</i>)

* Controls/Beheer = > 90% effective/doeltreffend

EFFICACY		DOELTREFFENDHEID
Milk tapeworm (<i>Moniezia expansa</i>)	CLASS1 / KLAS I	Melklintwurm (<i>Monezia expansa</i>)

Class I: 100 % effective in more than 80 % of treated animals.

Klas I: 100 % doeltreffend in meer as 80 % van behandelde diere.

EFFICACY		DOELTREFFENDHEID
Nasal bot (<i>Oestrus ovis</i>) larvae infestation	Controls / Beheer	Neuswurm (<i>Oestrus ovis</i>) larwe Infestasië

- †
- ② Benzimidazole
 - ⑧ Isoquinoline
 - ① Macrocyclic lactone

- †
- ② Bensimidasool
 - ⑧ Isokwinalien
 - ① Makrosikliese laktoon

Presentation

100 ml, 200 ml, 250 ml, 500 ml

Registration holder:

Afrivet Business Management (Pty)
Ltd Co. Reg. No. 2000/011263/07
P O Box 2009
Faerie Glen, 0043 R.S.A.
Tel. No. 012 817 9060
Web: www.afrivet.co.za
Helpline: 0860 833 2787

Aanbieding

100ml, 200 ml, 250 ml, 500 ml

Registrasiehouer:

Afrivet Business Management (Pty)
Ltd Mpy. Reg. No. 2000/011263/07
Posbus 2009
Faerie Glen, 0043 R.S.A.
Tel. No. 012 817 9060
Web: www.afrivet.co.za
Hulplyn: 0860 833 2787