

PACKAGE INSERT: ULTRATET

FOR ANIMAL USE ONLY

ULTRATET

Reg. No. G296 Act 36/1947; Namibia: V03.17.1.2/698
Zambia: 337/703V

STORAGE INSTRUCTIONS

Store in a cool, dark place.

COMPOSITION

ULTRATET is a stable, sterile solution containing 125 mg Oxytetracycline per ml.

INDICATIONS

For the treatment of tick-borne gallsickness (anaplasmosis), heartwater, pneumonia, footrot, joint-ill and navel-ill in cattle, sheep, goats and pigs, as well as strangles in horses.

WARNINGS

- DO NOT SLAUGHTER ANIMALS FOR HUMAN CONSUMPTION WITHIN 7 DAYS OF LAST TREATMENT.
- MILK AND MILK PRODUCTS FROM TREATED COWS MAY NOT BE USED FOR HUMAN CONSUMPTION WITHIN 48 HOURS OF LAST TREATMENT.
- Use only in cattle, sheep, goats, pigs and horses.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Intravenous injection should be done slowly into the chosen vein.
- At the first sign of discomfort, the injection should be stopped momentarily.
- Intramuscular injection can cause a slight transient discomfort, but may be minimized by massage of the injection site to disperse the drug.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

- Intramuscular
- Subcutaneous or
- SLOW INTRAVENOUS

N.B. Not more than 10 ml to be injection at any one site. Inject only into neck muscles in animals intended for slaughter for human consumption.

SLEGS VIR DIEREGERBUIK

ULTRATET

Reg. No. G296 Wet 36/1947; Namibia: V03.17.1.2/698
Zambia: 337/703V

BERGINGSAAWYSINGS

Berg in 'n koel, donker plek.

SAMESTELLING

ULTRATET is 'n stabiele, steriele oplossing wat 125 mg Oksitetrasiklien per ml bevat.

AANDUIDINGS

Vir die behandeling van bosluisoorgedraagde galsiekte (anaplasrose), hartwater, vrotpootjie, longontsteking, naelstringsiekte en septiese gewrigsontsteking by beeste, skape, bokke en varke asook nuwesiekte by perde.

WARNINGS

- MOENIE DIERE VIR MENSLIKE VERBRUIK BINNE 7 DAE NA LAASTE BEHANDELING SLAG NIE.
- MELK EN MELKPRODUKTE VAN BEHANDELDE DIERE MOENIE VIR MENSLIKE VERBRUIK BINNE 48 UUR NA DIE LAASTE BEHANDELING GEBRUIK WORD NIE.
- Gebruik slegs by beeste, skape, bokke, varke en perde.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

VOORSORGMATREËLS

- Binnearse inspuiting moet stadig in die gekose aar gegee word.
- By die eerste tekens van ongemak moet die inspuiting tydelike onderbreek word.
- Binnespiersse inspuiting kan 'n mate van verbygaande ongemak veroorsaak, maar deur die inspuitplek te masseer om die middel te versprei, kan dit beperk word.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

- Binnespiers
- Onderhuids of
- STADIG BINNEAARS

L.W. Daar moenie meer as 10 ml op een plek ingespuut word nie. Spuit net by die nekspiere in by diere wat vir menslike verbruik geslag gaan word.

Dosage/Dosis:

ANIMAL/DIER	INDICATIONS/AANWYSINGS	DOSAGE/DOSIS
Cattle/Beeste	Heartwater Tick-borne gallsickness (Anaplasmosis)	8 ml per 100 kg body mass daily as required
	Heartwater Bosluisoorgedraagde galsiekte (Anaplasrose)	8 ml per 100 kg liggaamsmassa daagliks soos benodig
	Footrot, Pneumonia, Navel-ill, Joint-ill	4 ml per 100 kg body mass daily for 3 – 4 days
	Vrotpootjie, Longontsteking. Naelstringsiekte, Septiese gewrigsonsteking	4 ml per 100 kg liggaamsmassa daagliks vir 3 – 4 dae
Horses/Perde	Strangles, Navel-ill, Joint-ill	4 ml per 100 kg body mass daily for 3 – 4 days
	Nuwesiekte, Naelstringsiekte. Septiese gewrigsonsteking	4 ml per 100 kg liggaamsmassa daagliks vir 3 – 4 dae
Sheep and goats/ Skape en bokke	Heartwater	4 ml per 50 kg body mass daily as required
	Heartwater	4 ml per 50 kg liggaamsmassa daagliks soos benodig
	Footrot, Pneumonia	2 ml per 50 kg body mass daily for 3 – 4 days
	Vrotpootjie, Longontsteking	2 ml per 50 kg liggaamsmassa daagliks vir 3 – 4 dae
Pigs/Varke	Footrot, Pneumonia	1 ml per 25 kg body mass daily for 3 – 4 days
	Vrotpootjie, Longontsteking	1 ml per 25 kg liggaamsmassa daagliks vir 3 – 4 dae

For rapid recovery a prompt and accurate diagnosis is essential. It is, therefore, always advisable to consult your veterinarian.

Vir vinnige herstel is 'n spoedige en akkurate diagnose noodsaaklik. Dit is dus altyd raadsaam om u veearts vir 'n korrekte diagnose te raadpleeg.

Presentation

50 ml, 100 ml, 500 ml

Aanbieding

50ml, 100 ml, 500 ml

Registration holder:

Afrivet Business Management (Pty) Ltd
Co. Reg. No. 2000/011263/07
P O Box 2009
Faerie Glen, 0043 R.S.A.
Tel. No. (012) 911-6416
Web: www.afrivet.co.za
Helpline: 0860 833 2787

Registrasiehouer:

Afrivet Business Management (Pty) Ltd
Mpy. Reg. No. 2000/011263/07
Posbus 2009
Faerie Glen, 0043 R.S.A.
Tel. No. (012) 911-6416
Web: www.afrivet.co.za
Hulplyn: 0860 833 2787