

WIREVAX PACKAGE INSERT/VOUBILJET

FOR ANIMAL USE ONLY

WIREVAX

Reg. No. G4200 Act 36/1947

STORAGE

Store in a cool place (2 °C – 8 °C).

Do not freeze.

Protect from light.

CONTAINS

Haemonchus contortus antigen 5 µg/ml

INDICATIONS

A vaccine to aid in the control of Wireworm infestations in sheep.

WARNINGS

- Do not mix with any other veterinary medicinal product.
- A transient swelling may appear at the site of inoculation for ± 13 to 17 days. If necessary consult a veterinarian.
- Animals may show a moderate rise of temperature for 1 to 3 days after vaccination. If necessary consult a veterinarian.
- Part used containers must be disposed of at the end of each day's operation since under field conditions it may be difficult to avoid accidental contamination of the vaccine. Burn or bury used containers.
- In any group of animals a small number of individuals may fail to respond to vaccination as a result of immunological incompetence. Satisfactory immune responses will only be attained in healthy animals, thus it is important to avoid vaccination of animals which have intercurrent infection or metabolic disorder. As with most killed vaccines, significant levels of immunity cannot be expected until two weeks after the primary vaccination course of three injections.
- Keep out of reach of children, uninformed persons and animals.
- Although this vaccine has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- In the case of accidental self-injection seek medical advice immediately and show the package leaflet to the physician.
- Only sterile needles and properly sterilized vaccinators should be used. No alcohol or other

SLEGS VIR DIEREGBRUIK

WIREVAX

Reg. No. G4200 Wet 36/1947

BERGING

Berg In 'n koelplek (2 °C – 8 °C).

Moenie bevries nie.

Beskerm teen lig.

BEVAT

Haemonchus contortus antigeen 5 µg/ml

AANDUIDINGS

'n Entstof wat help met die beheer van Haarwurmbesmetting in skape.

WAARSKUWINGS

- Moenie met 'n ander geneesmiddel meng nie.
- 'n Tydelike swelling by die inspuitplek mag voorkom vir ± 13 tot 17 dae. Indien nodig raadpleeg 'n veearts.
- Diere mag 'n matige verhoging in temperatuur vir 1 tot 3 dae toon. Indien nodig raadpleeg 'n veearts.
- Houers wat gedeeltelik gebruik is moet na daaglikse gebruik vernietig word aangesien dit onder veld toestande moeilik is om te verhoed dat die entstof besmet word. Verbrand of begrawe gebruikte houers.
- In 'n groep van diere mag 'n klein aantal individue versuim om te reageer op die inenting as 'n gevolg van immunologiese onbevoegdheid. Bevredigende immuunstatus sal slegs verkry word in gesonde diere. Dit is dus belangrik om nie diere te ent wat enige vorm van infeksie of metaboliese afwykings toon nie. Soos met die meeste dooie entstowwe, kan 'n beduidende vlak van immuniteit nie verwag word voordat twee weke verloop het na die eerste van die drie aanbevole inentings nie.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie entstof breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

VOORSORGMAATREËLS

- Indien die persoon wat die entstof toedien hom/haarsel per abuis inspuit, verkry onmiddellik mediese advies en toon die voubiljet aan die geneesheer.
- Slegs steriele naalde en behoorlik gesteriliseerde spuite moet gebruik word. Om higiëne te verseker is dit

disinfectants should be used for sterilisation. In order to maintain proper hygiene, it is advisable to change to clean sterile needles as frequently as possible.

- In case of long fleece, it is necessary to make sure that the needle has penetrated the skin before injecting the product. Use thin (20G) needles. Thick needles may cause leakage of the product.
- The injection site should be made through an area of clean dry skin.
- Avoid exposure to high temperatures and direct sunlight during inoculation.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

- MIX GENTLY BEFORE USE by inverting the bottle two or three times.
- May be used in pregnant animals.

DOSE AND ADMINISTRATION

Sheep of any age and lambs over 3 weeks

Use all product within 12 hours of opening.

- Inject 1 ml Wirevax subcutaneously, preferably using the inside of the thigh as the injection site.

EFFICACY

- To induce immunity, three injections are required at intervals of 3 to 4 weeks. For the remainder of the wireworm risk period, immunity can be maintained by boosters given at 6 week intervals.
- **It is important that sheep vaccinated with Wirevax do not share their pasture with unvaccinated animals of either species. Do not move vaccinated animals onto pasture recently grazed by unvaccinated sheep.**
- For protection of lambs born in September / October the following injection schedule is recommended for control of wireworm throughout the summer:

raadsaam om naalde gereeld te vervang met skoon steriele naalde.

- In die geval van langwol is dit noodsaaklik om seker te maak dat die naald die vel deurdring voor daar ingespuut word. Gebruik dun (20G) naalde. Dik naalde mag lekkasie van die produk veroorsaak.
- Die inspuitlek op die vel moet skoon en droog wees.
- Vermyn blootstelling aan hoë temperatuur en direkte sonlig gedurende inenting.

GEBRUIKSAANWYSINGS: GEBRUIK SELGS SOOS AANGEDUI

- SKUD SAGKENS VOOR GEBRUIK deur die flessie twee of drie keer heen en weer te kantel.
- Kan in dragtige diere gebruik word.

DOSIS EN TOEDIENING

Skape van enige ouderom en lammers ouer as 3 weke

- Gebruik binne 12 ure na opening.
- Spuit 1ml Wirevax onderhuids, verkieslik deur gebruik te maak van die binnekant van die lies as inspuitlek.

DOELTREFFENDHEID

- Drie inspuittings 3 tot 4 weke uitmekaar is nodig om beskerming te bewerkstellig. Vir die res van die haarwurm risikotydpark, kan immuniteit dan gehandhaaf word deur 6 weklike skraagdosisse.
- **Dit is belangrik dat skape wat met Wirevax ingeënt is, nie hulle weiveld deel met ongeënte diere nie. Moenie geënte diere op weiding plaas wat onlangs deur ongeënte skape beweë is nie.**
- Vir die beskerming van lammers wat gebore is in September / Oktober word die volgende entprogram aanbeveel vir die beheer van Haarwurm gedurende die somer:

Vaccination schedule/ Entprogram	± Lamb age (weeks)/ ± Lam ouderdom (weke)	Gap to next vaccination/ Intervalle tussen inertings	Time of year/ Tyd van die jaar	Could coincide with/ Kan saamval met
First/Eerste = V1	3 - 8	3 - 4 weeks/weke	Early November/ Vroeg November	Lamb marking/ Oormerking Castration or docking/ Kastrasie of stert afsit
Second/Tweede = V2	7 - 13	3 - 4 weeks/weke	Late November or early December Laat November of vroeg Desember	Weaning/Speen
Third/Derde = V3	11 - 17	6 weeks/weke	Late December or early Jan/ Laat Desember of vroeg Januarie	
Fourth/Vierde = V4	17 - 23	6 weeks/weke	6 weeks post V3 = February 6 weke na V3 = Februarie	Crutching/Mikskeer
Fifth/Vyfde = V5	23 - 29	Every 6 weeks up to 6 months/ Elke 6 weke tot op 6 maande	6 weeks post V4 = March or April 6 weke na V4 = Maart of April	

- If the spring is unusually wet, a dose of an effective anthelmintic is advised at the time of V2.
- This vaccination schedule can be used to control Wireworm in older animals.
- For lambs born in November or later, administration of an effective anthelmintic at the same time as V1 and V2 is recommended. Consult your veterinarian.
- Wirevax can be administered at the same time as a Clostridial vaccine though not at the same injection site.

- Indien die lente buitengewoon nat is, word dit aanbeveel dat 'n doeltreffende wurmmiddel toegedien word ten tye van die toediening van V2.
- Die entprogram kan gebruik word om Haarwurm in ouer diere te beheer.
- Vir lammers wat in November of later gebore word, word 'n doeltreffende wurmmiddel op dieselfde tyd as die V1 en V2 aanbeveel. Raadpleeg u veearts.
- Wirevax kan op dieselfde tyd as 'n Klostridium entstof toegedien word, maar nie op dieselfde inspuitplek nie.

Presentation

250 ml

Registration holder:

Afrivet Business Management (Pty) Ltd
Co. Reg. No. 2000/011263/07
P O Box 2009
Faerie Glen, 0043 R.S.A.
Tel: (012) 817 9060
Web: www.afrivet.co.za
Helpline: 0860 833 2787

Aanbieding

250 ml

Registrasiehouer:

Afrivet Business Management (Pty) Ltd
Mpy. Reg. No. 2000/011263/07
Posbus 2009
Faerie Glen, 0043 R.S.A.
Tel: (012) 817 9060
Web: www.afrivet.co.za
Hulplyn: 0860 833 2787